

Durham Research Online

Deposited in DRO:

08 March 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Müller, Juliane and Romero, Oscar and Cowan, Ellen A. and McClymont, Erin L. and Forwick, Matthias and Asahi, Hirofumi and März, Christian and Moy, Christopher M. and Suto, Itsuki and Mix, Alan and Stoner, Joseph (2018) 'Cordilleran ice-sheet growth fueled primary productivity in the Gulf of Alaska, northeast Pacific Ocean.', *Geology*, 46 (4). pp. 307-310.

Further information on publisher's website:

<https://doi.org/10.1130/G39904.1>

Publisher's copyright statement:

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.



UNIVERSITY OF LEEDS

This is a repository copy of *Cordilleran ice-sheet growth fueled primary productivity in the Gulf of Alaska, northeast Pacific Ocean*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/126648/>

Version: Accepted Version

Article:

Müller, J, Romero, O, Cowan, EA et al. (8 more authors) (2018) Cordilleran ice-sheet growth fueled primary productivity in the Gulf of Alaska, northeast Pacific Ocean. *Geology*, 46 (4). pp. 307-310. ISSN 0091-7613

<https://doi.org/10.1130/G39904.1>

© 2018 Geological Society of America. This is an author produced version of a paper published in *Geology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Cordilleran ice-sheet growth fueled primary productivity in the Gulf of Alaska, NE Pacific

Juliane Müller^{1,2}, Oscar Romero³, Ellen A. Cowan⁴, Erin L. McClymont⁵, Matthias Forwick⁶, Hirofumi Asahi⁷, Christian März⁸, Christopher M. Moy⁹, Itsuki Suto¹⁰, Alan Mix¹¹, Joseph Stoner¹¹

¹Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Am Alten Hafen 26, 27568 Bremerhaven, Germany

²Department of Geosciences, University of Bremen, Klagenfurter Straße 2-4, 28359 Bremen, Germany

³MARUM, Center for Marine Environmental Sciences, Leobener Straße 8, 28359 Bremen Germany

⁴Department of Geological and Environmental Sciences, Appalachian State University, ASU Box 32067, Boone, NC 28608-2067, USA

⁵Department of Geography, Durham University, Lower Mountjoy, South Road, Durham, DH1 3LE, UK

⁶Department of Geosciences, UiT The Arctic University of Norway in Tromsø, Postboks 6050 9037 Tromsø Langnes, Norway

⁷Korea Polar Research Institute, 12 Gaetbeol-ro, Yeonsu-gu, Incheon 406-840, South Korea

⁸School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

⁹Department of Geology, University of Otago, PO Box 56, Dunedin 9054, New Zealand

¹⁰Department of Earth and Planetary Sciences, Nagoya University, 464-8601 Furo-cho, Chikusa-ku, Nagoya, Japan

¹¹Oregon State University, 104 CEOAS Administration Building, Corvallis, OR 97331-5503,
USA

ABSTRACT

Fertilization of the ocean by eolian dust and icebergs is an effective mechanism to enhance primary productivity. In particular, high-nutrient, low-chlorophyll areas (HNLCs) where phytoplankton growth is critically iron (Fe)-limited, such as the subarctic Pacific and the Southern Ocean, are proposed to respond to increases in bioavailable Fe-supply with enhanced phytoplankton productivity and carbon export to the seafloor. While Fe-fertilization from dust is widely acknowledged to explain a higher export production during glacial periods in the Southern Ocean, paleoceanographic records supporting links between productivity and eolian dust and/or icebergs in the North Pacific are scarce. By combining independent proxies indicative of ice-sheet dynamics and ocean productivity from a single marine sedimentary record (IODP Site U1417), we present a comprehensive data set of phytoplankton response to different fertilization mechanisms in the subarctic northeast Pacific between 1.5 and 0.5 Ma, including the Mid Pleistocene Transition (MPT). Importantly, the timing of the fertilization events is more strongly controlled by local ice-sheet processes than by glacial-interglacial climate variability. Our findings indicate that fertilization by glacial debris results in productivity events in ocean areas adjacent to ice-sheets and that these mechanisms may represent an important, yet rarely considered driver of phytoplankton growth.

INTRODUCTION

The stimulation of primary productivity through the addition of Fe to the ocean surface, particularly in HNLC areas, significantly contributes to ocean carbon sequestration (Martin, 1990; Sigman et al., 2010). Field observations and laboratory experiments imply that, in addition to the input of Fe-rich eolian dust (Martin et al., 1989), delivery of macro- as well as

micronutrients and vertical mixing processes in the vicinity of icebergs foster phytoplankton growth in high latitude oceans (Duprat et al., 2016; Smith et al., 2007). Such in situ measurements and remote sensing data suggest a potentially important role for icebergs and eolian dust in driving primary productivity in HNLC regions, but provide only a snapshot view of modern ocean biogeochemical feedbacks. Paleoreconstructions, in turn, permit an integrated view and evaluation of the role of these fertilization mechanisms on export production. Owing to its proximity to a former major Northern Hemisphere ice-sheet, the Gulf of Alaska (GoA; NE Pacific) is an area with vigorous temperate glacial erosion of Fe-rich rocks (Gulick et al., 2015; Montelli et al., 2017). Here, we present the first reconstruction of phytoplankton productivity in the GoA linked to Fe inputs from glacial debris. We focus on sediments spanning the last important climate transition in Earth's history, the Mid Pleistocene Transition (MPT), when the Northern Cordilleran Ice Sheet (NCIS) experienced a significant expansion (Gulick et al., 2015). Although the exact timing and cause(s) of the MPT are intensely discussed (Clark et al., 2006; Elderfield et al., 2012; Maslin and Brierley, 2015), the potential for biogeochemical feedbacks operating in the high-latitude oceans during this crucial time interval of northern hemisphere ice-sheet growth remains poorly studied. This is the first assessment of (subpolar) Fe-fertilization mechanisms across the MPT from outside the Southern Ocean (Lamy et al., 2014; Martinez-Garcia et al., 2011).

We present a multi-proxy record including geochemical, micropaleontological and sedimentological data obtained from IODP Site U1417 in the GoA (56°57'N, 147°6'W, 4200 m water depth; DR1; Jaeger et al., 2014). Our results record the interactions between sea surface temperature (SST), the input of terrigenous material by both eolian as well as ice rafting processes, and export productivity for multiple glacial-interglacial cycles between 1.5 and 0.5 Ma (Fig. 1). In the absence of eolian dust measurements, elevated contents of land-plant specific long-chain n-alkanes (depicted by higher terrigenous-aquatic ratios (TAR); Meyers, 1997; Peters et al., 2004) are used to track terrestrial dust input (Simoneit, 1977). In addition, icebergs

may carry high amounts of terrigenous organic matter to distal ocean sites and are considered as a further transport agent of these leaf-wax compounds (Knies, 2005; Stein et al., 2009; Villanueva et al., 1997). Accordingly, at Site U1417, elevated TAR values that coincide with at ice-rafted debris (IRD) maxima suggest an ice rafting of leaf-wax lipids, while maximum TAR values accompanied by IRD minima indicate an airborne transport of these compounds. From the consistent pattern in concurrently high marine productivity indicators and high TAR values, we deduce that enhanced marine productivity was directly related to the input of terrigenous matter. Details on individual analytical methods and the age model are provided as Supplementary Information DR2.

Sea surface conditions and different Fe-fertilization mechanisms in the GoA

An overall consistent relationship applies at U1417, with intervals of lower SSTs and more polar waters (%C_{37:4}) coinciding with higher deposition of IRD (e.g., MIS 39, 30, 20), indicating a direct link between GoA sea surface conditions and NCIS dynamics. A distinct variability in diatom abundances, biogenic silica (opal; BSi) content and the Ba/Al ratio is considered to reflect abrupt phytoplankton productivity changes at Site U1417 (Fig. 1). Despite relatively warm SSTs prior to the MPT (> 1.2 Ma), the occurrence of diatoms was confined to short-lived events, and a significant rise in diatom abundance and biogenic silica content occurred only at the onset of the MPT (1.22 Ma, MIS 37; Fig. 1). The association between the biosiliceous signal and SST is not consistent over the entire record and SST changes do not appear to be a primary driver of diatom productivity. However, both diatom and BSi signals are strongly linked to elevated Ba/Al values, recording increased export productivity (Jaccard et al., 2010), and to higher TAR values (Fig. 1). Today, significant amounts of Fe-rich glacial silt are deposited along glaci-fluvial river banks and at glacier termini along South Alaskan coastal areas and glacial rock flour is transported beyond the continental shelf into Fe-limited pelagic waters during dust storms (Crusius et al., 2011; Muhs et al., 2016). Evidently, the eolian

transport of this glacial flour- derived dust via strong northerly winds is an important mechanism for the supply of bioavailable Fe to foster phytoplankton blooms in the GoA (Crusius et al., 2011; Crusius et al., 2017). We hence argue that the TAR peaks coinciding with diatom, BSi and Ba/Al maxima and IRD minima at Site U1417 reflect intervals of enhanced eolian export of leaf-wax lipids together with Fe-rich Alaskan dust, leading to productivity increases in the GoA across the MPT (e.g., at 1.22, 1.15 and 0.99 Ma; Fig. 1; DR3). Similarly, McDonald et al. (1999) proposed that late Pleistocene diatom productivity events at ODP Site 887 could have been promoted by Fe-supply via dust.

In addition to dust-fertilization, we suggest that also ice rafting of glacial Fe-rich debris (transported together with glacially reworked organic matter containing leaf-wax lipids) stimulated productivity at Site U1417. Intervals characterised by enhanced IRD deposition and high TAR, diatom, BSi and Ba/Al values occurred at e.g. 1.05, 0.91, 0.77 and 0.66 Ma (Fig. 1; DR3). Recent observations highlight the importance of Fe-fertilization of pelagic ecosystems from icebergs, accounting for up to 20% of the total carbon export in the Southern Ocean (Duprat et al., 2016; Smith et al., 2007). The coincidence of ice rafting and elevated marine productivity events in the GoA suggests that this mechanism also operated during the MPT in the subpolar NE Pacific. In addition to dust- and iceberg-fertilization, Fe-supply via mesoscale eddies (Crawford et al., 2007) and volcanic ash (Hamme et al., 2010) may have promoted phytoplankton blooms in the GoA. However, we consider these mechanisms of only minor importance at Site U1417 (see DR4 for discussion).

From the early (> 1 Ma) towards the late (> 0.6 Ma) MPT, we note a decrease in predominantly dust-fertilized productivity pulses, while iceberg-fertilization sustained. This transition could result from an overall reduction in dust export owing to the persistent expansion of the NCIS (sealing central Alaskan dust (loess) deposits) and/or a change in atmospheric circulation diverting Alaskan storm tracks. Deposition of lithic particles by ice rafting, however, does not per se relate to a higher export production in the GoA and we argue that additional factors

impacted ocean productivity (e.g. nitrate depletion; Galbraith et al., (2008)). Peaks in IRD at 1.27 or 0.82 Ma, for example, do not coincide with higher Ba/Al or opal values but an enhanced abundance of the C_{37:4} alkenone (Fig. 1), pointing to a significantly cooler ocean surface.

Further implications

With regard to the overall environmental evolution in the subpolar NE Pacific, we suggest that the diatom and opal peaks at 1.22 Ma mark a transition when NCIS growth and, hence, the production and export of glaciogenic dust led to an effective Fe-fertilization in the adjacent GoA. Whereas eolian dust-fertilization dominated during intervals of reduced glacier extent (i.e., when coastal plains and glaciogenic silt deposits were subaerially exposed; Fig. 2A, B), iceberg-fertilization occurred during intervals of enhanced glaciation when the NCIS terminated on the Alaskan continental shelf and discharged icebergs to Site U1417 (Fig. 2C, D). We note that, during the latter intervals, strong katabatic winds may have sustained an (airborne) export of dust from areas that remained ice-free (DR3).

Interestingly, the higher dust input at Site U1417 at approximately 1.22 Ma coincides with an enormous increase in dust delivery to the subantarctic Atlantic (Martinez-Garcia et al., 2011). Ocean cooling as well as increasing latitudinal temperature gradients are considered to have accounted for an equatorward movement of oceanic fronts and a strengthened atmospheric circulation leading to a higher dust export to the subantarctic Southern Ocean during the MPT (Kemp et al., 2010; Martinez-Garcia et al., 2011; McClymont et al., 2013). We suggest that the expansion of polar waters in the high northern latitudes and the growth of the NCIS (affecting surface albedo and orography) could have induced similar atmospheric shifts promoting dust export events in the GoA at the onset of the MPT. Comparisons between northwestern and eastern records of subpolar North Pacific paleoproductivity, however, reveal that although SSTs in both areas developed in a similar fashion, the patterns of Mid Pleistocene primary productivity did not. While export production generally decreased in the Bering Sea due to an

155 increase in sea ice cover (Kim et al., 2014), the productivity events observed in the GoA point
156 to an efficient, yet sporadic, ocean fertilization from the input of NCIS-sourced glacigenic
157 terrestrial matter (and Fe) across the MPT.

158 We note that the productivity pulses at Site U1417 are neither exclusively confined to glacials
159 nor to interglacials. This pattern contrasts to the western subarctic Pacific and the Bering Sea,
160 where opal production increased primarily during Pleistocene interglacials (Kim et al., 2014).

161 The productivity pulses at Site U1417 may reflect local feedback mechanisms between South
162 Alaskan glacier dynamics (controlling ice-proximal dust production and dispersal), and an
163 immediate response of the marine ecosystem, yet they highlight potentially relevant
164 mechanisms to elucidate hitherto neglected interactions in the land-ocean-atmosphere system
165 during glacial-interglacial transitions. We propose the GoA as a case example of a Pleistocene
166 ice-proximal marine environment where ice-sheet dynamics exhibited a significant control on
167 primary productivity and potentially also CO₂ draw-down. In fact, with the intensification of
168 Pleistocene Northern Hemisphere glaciation and sea-level lowering, extensive sub-aerial pro-
169 glacial (coastal) outwash plains developed not only in South Alaska but also along the
170 Laurentide Ice Sheet and European Ice Sheets, and these areas should be considered as
171 potentially important sources of Fe-bearing glacigenic silt (Bullard et al., 2016) for areas where
172 seasonal Fe-limitation restricts phytoplankton growth (Moore et al., 2006; Nielsdóttir et al.,
173 2009). Further exploration of sedimentary archives from high-latitude ocean areas adjacent to
174 (paleo) ice-sheets that permit correlations between productivity proxies and terrigenous
175 compounds are required to evaluate the potential impacts of glacigenic dust- and iceberg-
176 fertilization on phytoplankton productivity across the MPT and beyond. Importantly, such data
177 would provide for a quantitative assessment of whether these processes could have accounted
178 for an amplification of glacial-interglacial cycles, or if they even contributed to an appreciable
179 CO₂ draw-down during the MPT.

We thank the IODP-USIO and the captain and crew of the D/V JOIDES Resolution. Funding was provided by the German Research Foundation (MU3670/1-2), an ECORD Research Grant and Helmholtz Association Grant VH-NG 1101, from NERC (IODP Rapid Response Award, NE/L002426/1) and a Philip Leverhulme Prize, from U.S. NSF award OCE-1434945 and post-expedition award from the U.S. Science Support Program of IODP, from Korea Polar Research Institute's Basic Research Project (PE16062) and a National Research Foundation of Korea Grant funded by the Korean Government (2015M1A5A1037243), from IODP After Cruise Research Program from JAMSTEC (H28-01), from JSPS KAKENHI Grant (JP26281006). This is a contribution to the AWI Helmholtz Research Programme PACES II WP3.1.

References

- Bullard, J. E., Baddock, M., Bradwell, T., Crusius, J., Darlington, E., Gaiero, D., Gassó, S., Gisladottir, G., Hodgkins, R., McCulloch, R., McKenna-Neuman, C., Mockford, T., Stewart, H., and Thorsteinsson, T., 2016, High-latitude dust in the Earth system: Reviews of Geophysics, v. 54, no. 2, p. 447-485.
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G., and Roy, M., 2006, The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂: Quaternary Science Reviews, v. 25, no. 23–24, p. 3150-3184.
- Crawford, W. R., Brickley, P. J., and Thomas, A. C., 2007, Mesoscale eddies dominate surface phytoplankton in northern Gulf of Alaska: Progress in Oceanography, v. 75, no. 2, p. 287-303.
- Crusius, J., Schroth, A. W., Gassó, S., Moy, C. M., Levy, R. C., and Gatica, M., 2011, Glacial flour dust storms in the Gulf of Alaska: Hydrologic and meteorological controls and

206 their importance as a source of bioavailable iron: *Geophysical Research Letters*, v.
207 38, no. 6, p. L06602.

208 Crusius, J., Schroth, A. W., Resing, J. A., Cullen, J., and Campbell, R. W., 2017, Seasonal and
209 spatial variabilities in northern Gulf of Alaska surface water iron concentrations
210 driven by shelf sediment resuspension, glacial meltwater, a Yakutat eddy, and dust:
211 *Global Biogeochemical Cycles*, v. 31, no. 6, p. 942-960.

212 Duprat, L. P. A. M., Bigg, G. R., and Wilton, D. J., 2016, Enhanced Southern Ocean marine
213 productivity due to fertilization by giant icebergs: *Nature Geosci*, v. 9, no. 3, p. 219-
214 221.

215 Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., McCave, I. N., Hodell, D., and
216 Piotrowski, A. M., 2012, Evolution of Ocean Temperature and Ice Volume Through
217 the Mid-Pleistocene Climate Transition: *Science*, v. 337, no. 6095, p. 704-709.

218 Galbraith, E. D., Kienast, M., Jaccard, S. L., Pedersen, T. F., Brunelle, B. G., Sigman, D. M.,
219 and Kiefer, T., 2008, Consistent relationship between global climate and surface
220 nitrate utilization in the western subarctic Pacific throughout the last 500 ka:
221 *Paleoceanography*, v. 23, no. 2, p. n/a-n/a.

222 Gulick, S. P. S., Jaeger, J. M., Mix, A. C., Asahi, H., Bahlburg, H., Belanger, C. L., Berbel, G.
223 B. B., Childress, L., Cowan, E., Drab, L., Forwick, M., Fukumura, A., Ge, S., Gupta,
224 S., Kioka, A., Konno, S., LeVay, L. J., März, C., Matsuzaki, K. M., McClymont, E.
225 L., Moy, C., Müller, J., Nakamura, A., Ojima, T., Ribeiro, F. R., Ridgway, K. D.,
226 Romero, O. E., Slagle, A. L., Stoner, J. S., St-Onge, G., Suto, I., Walczak, M. D.,
227 Worthington, L. L., Bailey, I., Enkelmann, E., Reece, R., and Swartz, J. M., 2015,
228 Mid-Pleistocene climate transition drives net mass loss from rapidly uplifting St.
229 Elias Mountains, Alaska: *Proceedings of the National Academy of Sciences*, v. 112,
230 no. 49, p. 15042-15047.

231 Hamme, R. C., Webley, P. W., Crawford, W. R., Whitney, F. A., DeGrandpre, M. D., Emerson,
 232 S. R., Eriksen, C. C., Giesbrecht, K. E., Gower, J. F. R., Kavanaugh, M. T., Peña,
 233 M. A., Sabine, C. L., Batten, S. D., Coogan, L. A., Grundle, D. S., and Lockwood,
 234 D., 2010, Volcanic ash fuels anomalous plankton bloom in subarctic northeast
 235 Pacific: *Geophysical Research Letters*, v. 37, no. 19.

236 Jaccard, S. L., Galbraith, E. D., Sigman, D. M., and Haug, G. H., 2010, A pervasive link
 237 between Antarctic ice core and subarctic Pacific sediment records over the past 800
 238 kyrs: *Quaternary Science Reviews*, v. 29, no. 1–2, p. 206-212.

239 Jaeger, J., Gulick, S. P. S., LeVay, L., Asahi, H., Bahlburg, H., Belanger, C., Berbel, G. B. B.,
 240 Childress, L., Cowan, E., Davies, M. H., Drab, L., Dottori, F., Forwick, M.,
 241 Fukumura, A., Ge, S., Gupta, S., Kioka, A., Konno, S., März, C., Matsuzaki, K.,
 242 McClymont, E., Mix, A., Moy, C., Müller, J., Nakamura, A., Ojima, T., Ridgway,
 243 K., Romero, O., Slagle A., Stoner, J., St-Onge, G., Suto, I., and Worthington, L.,
 244 2014, Southern Alaska margin: Interactions of tectonics, climate, and
 245 sedimentation., *Proceedings of the Integrated Ocean Drilling Program*, vol. 341,
 246 doi:10.2204/iodp.proc.341.101.2014.

247 Kaufman, D. S., Young, N. E., Briner, J. P., and Manley, W. F., 2011, Chapter 33 - Alaska
 248 Palaeo-Glacier Atlas (Version 2), in Jürgen Ehlers, P. L. G., and Philip, D. H., eds.,
 249 *Developments in Quaternary Sciences*, Volume 15, Elsevier, p. 427-445.

250 Kemp, A. E. S., Grigorov, I., Pearce, R. B., and Naveira Garabato, A. C., 2010, Migration of
 251 the Antarctic Polar Front through the mid-Pleistocene transition: evidence and
 252 climatic implications: *Quaternary Science Reviews*, v. 29, no. 17–18, p. 1993-2009.

253 Kim, S., Takahashi, K., Khim, B.-K., Kanematsu, Y., Asahi, H., and Ravelo, A. C., 2014,
 254 Biogenic opal production changes during the Mid-Pleistocene Transition in the
 255 Bering Sea (IODP Expedition 323 Site U1343): *Quaternary Research*, v. 81, no. 1,
 256 p. 151-157.

Knies, J., 2005, Climate-induced changes in sedimentary regimes for organic matter supply on the continental shelf off northern Norway: *Geochimica et Cosmochimica Acta*, v. 69, no. 19, p. 4631-4647.

Lamy, F., Gersonde, R., Winckler, G., Esper, O., Jaeschke, A., Kuhn, G., Ullermann, J., Martinez-Garcia, A., Lambert, F., and Kilian, R., 2014, Increased Dust Deposition in the Pacific Southern Ocean During Glacial Periods: *Science*, v. 343, no. 6169, p. 403-407.

Lisiecki, L. E., and Raymo, M. E., 2005, A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records: *Paleoceanography*, v. 20, no. 1, p. PA1003.

Martin, J. H., 1990, Glacial-interglacial CO_2 change: The Iron Hypothesis: *Paleoceanography*, v. 5, no. 1, p. 1-13.

Martin, J. H., Gordon, R. M., Fitzwater, S., and Broenkow, W. W., 1989, Vertex: phytoplankton/iron studies in the Gulf of Alaska: *Deep Sea Research Part A. Oceanographic Research Papers*, v. 36, no. 5, p. 649-680.

Martinez-Garcia, A., Rosell-Mele, A., Jaccard, S. L., Geibert, W., Sigman, D. M., and Haug, G. H., 2011, Southern Ocean dust-climate coupling over the past four million years: *Nature*, v. 476, no. 7360, p. 312-315.

Maslin, M. A., and Brierley, C. M., 2015, The role of orbital forcing in the Early Middle Pleistocene Transition: *Quaternary International*, v. 389, p. 47-55.

McClymont, E. L., Soudian, S. M., Rosell-Melé, A., and Rosenthal, Y., 2013, Pleistocene sea-surface temperature evolution: Early cooling, delayed glacial intensification, and implications for the mid-Pleistocene climate transition: *Earth-Science Reviews*, v. 123, p. 173-193.

McDonald, D., Pedersen, T. F., and Crusius, J., 1999, Multiple late Quaternary episodes of exceptional diatom production in the Gulf of Alaska: *Deep Sea Research Part II: Topical Studies in Oceanography*, v. 46, no. 11-12, p. 2993-3017.

283 Montelli, A., Gulick, S. P. S., Worthington, L. L., Mix, A., Davies-Walczak, M., Zellers, S. D.,
 284 and Jaeger, J. M., 2017, Late Quaternary glacial dynamics and sedimentation
 285 variability in the Bering Trough, Gulf of Alaska: *Geology*, v. 45, no. 3, p. 251-254.
 286 Moore, C. M., Mills, M. M., Milne, A., Langlois, R., Achterberg, E. P., Lochte, K., Geider, R.
 287 J., and La Roche, J., 2006, Iron limits primary productivity during spring bloom
 288 development in the central North Atlantic: *Global Change Biology*, v. 12, no. 4, p.
 289 626-634.
 290 Muhs, D. R., Budahn, J. R., Skipp, G. L., and McGeehin, J. P., 2016, Geochemical evidence
 291 for seasonal controls on the transportation of Holocene loess, Matanuska Valley,
 292 southern Alaska, USA: *Aeolian Research*, v. 21, p. 61-73.
 293 Nielsdóttir, M. C., Moore, C. M., Sanders, R., Hinz, D. J., and Achterberg, E. P., 2009, Iron
 294 limitation of the postbloom phytoplankton communities in the Iceland Basin: *Global*
 295 *Biogeochemical Cycles*, v. 23, no. GB3001.
 296 Sigman, D. M., Hain, M. P., and Haug, G. H., 2010, The polar ocean and glacial cycles in
 297 atmospheric CO₂ concentration: *Nature*, v. 466, no. 7302, p. 47-55.
 298 Simoneit, B. R. T., 1977, Organic matter in eolian dusts over the Atlantic Ocean: *Marine*
 299 *Chemistry*, v. 5, no. 4, p. 443-464.
 300 Smith, K. L., Robison, B. H., Helly, J. J., Kaufmann, R. S., Ruhl, H. A., Shaw, T. J., Twining,
 301 B. S., and Vernet, M., 2007, Free-Drifting Icebergs: Hot Spots of Chemical and
 302 Biological Enrichment in the Weddell Sea: *Science*, v. 317, no. 5837, p. 478-482.
 303 Stein, R., Hefter, J., Grützner, J., Voelker, A., and Naafs, B. D. A., 2009, Variability of surface
 304 water characteristics and Heinrich-like events in the Pleistocene midlatitude North
 305 Atlantic Ocean: Biomarker and XRD records from IODP Site U1313 (MIS 16-9):
 306 *Paleoceanography*, v. 24, p. PA2203.


307 Villanueva, J., Grimalt, J. O., Cortijo, E., Vidal, L., and Labeyrie, L., 1997, A biomarker
308 approach to the organic matter deposited in the North Atlantic during the last
309 climatic cycle: *Geochimica et Cosmochimica Acta*, v. 61, no. 21, p. 4633-4646.

310

FIGURE CAPTIONS

Figure 1: Records of phytoplankton productivity (diatom concentration, BSi content, Ba/Al), terrigenous-aquatic ratio (TAR), IRD (3-point running average of wt.% coarse sand grains) deposition, and SST (U_{37}^K , $U_{37}^{K'}$, %C_{37:4}) at Site U1417 compared to the $\delta^{18}O$ isotope stack (Lisiecki and Raymo, 2005) over 1.5 - 0.5 Ma. Blue shadings highlight glacial intervals. Filled and hollow circles mark high productivity events stimulated by iceberg- and eolian dust-fertilization, respectively. Gray numbers mark Marine Isotope Stages (MIS).

Figure 2: Site U1417 (56°57'N, 147°6'W) and different Mid Pleistocene environmental settings in the study area and associated fertilization mechanisms. Brown shadings refer to modern Alaskan loess deposits (after Muhs et al., 2016). A, B: Reduced ice-sheet coverage (pale blue shadings) and a predominantly eolian export of glacial dust to Site U1417. C, D: Periods of an extended NCIS (2C; after Kaufman et al., 2011) with marine terminating glaciers and ice-rafting of glacial debris across the GoA. Green shadings indicate assumed area of dust- and iceberg-fertilized high productivity in the GoA through the MPT.



328

329

